



Health and
Wellness

Prince Edward Island Guidelines for the Management and Control of *Verotoxic E Coli*

January 2018

Department of Health and Wellness
Chief Public Health Office

Verotoxic *E coli*

Contents

Case Definition ¹	2
Reporting Requirements.....	2
Etiology	2
Clinical Presentation	2
Diagnosis	3
Epidemiology ³	3
Occurrence.....	3
Control	4
References	7

Verotoxic *E coli*

Case Definition¹

Confirmed Case¹

Laboratory confirmation of infection with or without clinical illness ^[1]:

- Isolation of verotoxin-producing *Escherichia coli* (includes but not limited to O157:H7) from an appropriate clinical specimen (e.g., feces, urine, blood)

OR

- Detection of verotoxin antigen or nucleic acid.

Probable Case

Clinical illness ^[1] in a person epidemiologically linked to a confirmed case which would include persons with haemolytic uremic syndrome (HUS).

^[1] Clinical illness is characterized by diarrhea (often bloody) and abdominal cramps, fever is often absent. Illness may be complicated by HUS, thrombocytopenia purpura (TTP) or pulmonary edema. Asymptomatic infections may also occur and the microorganism may cause extra-intestinal infections.

Reporting Requirements

Laboratories

The Provincial Laboratory shall in accordance with the Prince Edward Island *Public Health Act*², report all positive laboratory results by phone and mail, fax or electronic transfer as soon as the result is known to the Chief Public Health Officer (CPHO) (or designate).

Etiology

Escherichia coli are gram-negative bacilli. The bacteria cause illness by creating a toxin referred to as a verotoxin (VTEC) or shiga-like toxin (STEC). The organism has a low infective dose.

Clinical Presentation

Similar to other foodborne illnesses, the symptoms of verotoxin-producing *Escherichia coli* (VTEC) involve the gastrointestinal tract including: acute diarrhea, cramps, nausea, emesis and occasional fever. Diarrhea can range from mild and non-bloody to stools that are virtually all blood. The illness is often self-limited. Most individuals recover in 5 to 10 days.

Approximately 5-10% of those with VTEC can develop HUS which develops when the verotoxin breaks down the lining of the intestines and in some cases, damages the kidneys. This occurs in up to 15% of children and the elderly. The overall case fatality of VTEC is 1%.

Diagnosis

The diagnosis is made by positive stool, urine and blood culture for *E. coli* O157:H7. However, other serotypes of *E.coli* have also produced verotoxin such as O145. The diagnosis of VTEC should be considered in the presence of severe diarrhea, HUS, TTP or hemorrhagic colitis.

Epidemiology³

1. Reservoir

Cattle are the principle reservoir of *Escherichia coli* including O157:H7. Humans may serve as an accidental host and therefore, act as a reservoir for person to person transmission.

2. Transmission

The predominant mode of transmission is through the ingestion of contaminated food, often related to inadequate cooking or through cross-contamination during food preparation. Foods (in particular fruits and vegetables) contaminated by cow manure have been documented modes of transmission. Serious outbreaks have occurred in North America from inadequately cooked hamburgers, unpasteurized milk or cheese, apple cider, alfalfa sprouts, dry-cured salami, lettuce, game meat, and cheese curds. Transmission also occurs directly from person to person (in families, daycares, and institutions). Infection can occur after swimming in or drinking contaminated water. Outbreaks in children have been associated with petting zoos.

3. Incubation Period

The incubation period is typically three to four days with a range of one to 10 days.

4. Period of Communicability

The infection is communicable for the duration of excretion of the pathogen, commonly a week or less in adults and up to approximately three weeks in about one-third of children. Prolonged carriage is uncommon.

5. Host susceptibility

The elderly and children appear to be at higher risk for illness. Children less than five years of age are at greatest risk of developing HUS.

Occurrence

1. General

VTEC is an important problem in North America, Europe, South Africa, Japan, the southern cone of South America, and Australia.

2. Canada

Sporadic cases and outbreaks have occurred in Canada since the first reported outbreak in the United States in 1982. The highest age specific rate is among children one to four years old. Overall, the highest incidence is found in children under the age of 15 years⁴.

3. Prince Edward Island

In the past ten years, the number of cases has ranged from two cases up to 16⁵.

Control

1. Management of a case

- The CPHO is involved with the investigation of all VTEC cases. Public Health Nursing, Health PEI, will follow up all lab confirmed cases and environmental health officers may be consulted on cases as appropriate. Advice on the management of cases will be provided by the CPHO.
- Notification of test results and prescription of treatment (if required) will be carried out by the attending health care provider.
- Information should be provided about disease transmission and the appropriate infection prevention and control measures to be implemented to minimize the possibility of transmission including strict hand hygiene especially after using the washroom, changing diapers and before preparing/handling and serving food.
- Provide direction on exclusion from work and day care/school. Exclusion is required for symptomatic persons who are:
 - food handlers whose work involves
 - touching unwrapped food to be consumed raw or without further cooking and/or
 - handling equipment or utensils that touch unwrapped food to be consumed raw or without further cooking,
 - healthcare, daycare or other staff who have contact through serving food with highly susceptible patients or persons, in whom an intestinal infection would have particularly serious consequences,
 - those involved in patient care or care of young children, elderly or dependent persons,
 - children attending daycares or similar facilities who are diapered or unable to implement good standards of personal hygiene, and
 - older children or adults who are unable to implement good standards of personal hygiene (e.g., mentally or physically challenged).

Verotoxic *E coli*

- Exclusion applies until **two stool specimens** taken from the case are reported as negative. These specimens should not be taken less than 24 hours apart and should be taken at least 48 hours after normal stools have resumed (lab requisition should note “test for clearance”).
- Advise the case about proper food handling practices, and to refrain from preparing food for others for the duration of the period of communicability.
- Contact precautions should be used in healthcare settings where children or adults have poor hygiene or incontinence that cannot be contained. Otherwise, routine practices are adequate.

2. Treatment of a case

- The use of antibiotics is not recommended and may be harmful by enhancing the release of toxins.
- Antimotility agents should be avoided.
- Replace fluids and electrolytes as required.

3. Management of contacts

- Contacts should be instructed about disease transmission, appropriate personal hygiene, routine practices, and contact precautions.
- Symptomatic contacts
 - Contacts who are symptomatic should be assessed by a physician.
 - Contacts who are symptomatic must be excluded from daycare or similar facilities or occupations involving food handling, patient care or care of young, elderly or dependent persons until they are no longer symptomatic.
- One stool specimen or culture will be requested from symptomatic contacts and must be reported as negative prior to returning to daycare or similar facilities, or occupations involving food handling, patient care or care of young, elderly or dependent persons.
- Asymptomatic contacts
 - Cultures of asymptomatic contacts should be done in consultation with the CPHO and would generally be confined to food handlers, attendants, and children in child care centers and other settings where the spread of infection is particularly likely.
 - The specimen must be reported as negative prior to returning to daycare or similar facilities or occupations involving food handling, patient care or care of young, elderly or dependent persons.

4. Preventative measures

- Provide public education about personal hygiene, especially the sanitary disposal of feces and careful hand washing after defecation and sexual contact, and before preparing or eating food.
- Advise infected individuals to avoid food preparation.
- Educate food handlers about proper food and equipment handling, preparation, and hygiene especially in avoiding cross-contamination from raw meat products, and thorough hand washing.
- Advise individuals to consume only pasteurized milk, dairy products, and juices.
- Advise on safe food preparation such as cooking beef adequately, especially ground beef, to an internal temperature of 71⁰C (160⁰F) and using a meat thermometer and not relying on cooking until pink color is gone; washing fruits and vegetables, particularly if eaten raw; and peeling raw fruits when possible.
- Educate about the risk of sexual practices that permit fecal-oral contact.
- Advise on testing private water supplies for presence of bacterial contamination, if suspected.
- Advise on washing hands thoroughly and frequently using soap, in particular after contact with farm animals or the farm environment, contact with animals in public settings such as fairs, farm tours, petting zoos, camps, and schools.

References

1. [Case Definitions for Communicable Diseases](#). *Public Health Agency of Canada*. [Online] November 2009.
2. Province of PEI. Public Health Act R.S.P.E.I [Internet]. 2013. Available from: http://www.gov.pe.ca/law/statutes/pdf/p-30_1.pdf
3. Heymann, David L. 2015. *Control of Communicable Diseases Manual 20th Edition*. Washington : American Public Health Association, 2015.
4. National Enteric Surveillance Program (NESP). [Annual Summary 2013](#). *Public Health Agency of Canada*.
5. CPHO Report 2016. *Health for All Islanders, Promote, Prevent, Protect: PEI Chief Public Health Officer's Report 2016*
6. Government of Alberta. *Escherichia coli Verotoxigenic Infections Guideline 2011*. [Online] August 2017. <https://open.alberta.ca/publications/escherichia-coli-verotoxigenic-infections>